

## ATION PAGE

Form Approved  
OMB No. 0704-01881a. REPORT  
Uncla

AD-A225 869

2a. SECURITY  
N/A2b. DECLASSIFICATION/DOWNGRADING SCHEDULE  
N/A

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

Technical Report No. 299

6a. NAME OF PERFORMING ORGANIZATION  
University of North Carolina  
Center for Stochastic Processes6b. OFFICE SYMBOL  
(If applicable)

1b. RESTRICTIVE MARKINGS

3. DISTRIBUTION/AVAILABILITY OF REPORT

Approved for Public Release;  
Distribution Unlimited

5. MONITORING ORGANIZATION REPORT NUMBER(S)

AFOSR-TR- 90 0854

7a. NAME OF MONITORING ORGANIZATION

AFOSR/NM

6c. ADDRESS (City, State, and ZIP Code)  
Statistics Department  
CB #3260, Phillips Hall  
Chapel Hill, NC 27599-3260

7b. ADDRESS (City, State, and ZIP Code)

Bldg. 410  
Bolling Air Force Base, DC 20332-64488a. NAME OF FUNDING/SPONSORING  
ORGANIZATION  
AFOSR8b. OFFICE SYMBOL  
(If applicable)  
NM

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

F49620 85C 0144

8c. ADDRESS (City, State, and ZIP Code)

Bldg. 410  
Bolling AFB, DC 20332-6448

10. SOURCE OF FUNDING NUMBERS

PROGRAM  
ELEMENT NO.  
6.1102FPROJECT  
NO.  
2304TASK  
NO.  
A5WORK UNIT  
ACCESSION NO.

11. TITLE (Include Security Classification)

On the upper and lower classes for stationary Gaussian random fields on Abelian groups  
with a regularly varying entropy

12. PERSONAL AUTHOR(S)

Albin, J.M.P.

13a. TYPE OF REPORT

preprint

13b. TIME COVERED

FROM \_\_\_\_\_ TO \_\_\_\_\_

14. DATE OF REPORT (Year, Month, Day)

1990, June

15. PAGE COUNT

12

16. SUPPLEMENTARY NOTATION

None.

17. COSATI CODES

FIELD GROUP SUB-GROUP

XXXXXXXXXXXXXXXXXXXX

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Upper and lower classes, LIL, Gaussian random fields.

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

We give a complete and relatively explicit characterization of the upper  
and lower classes for a general stationary Gaussian random field.DTIC  
ELECTE  
AUG 29 1990  
S D

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

22a. NAME OF RESPONSIBLE INDIVIDUAL

Professor Eytan Barouch Jon A. Sjogren

22b. TELEPHONE (Include Area Code)


(202) 767-5026

22c. OFFICE SYMBOL

AFOSR/NM

# CENTER FOR STOCHASTIC PROCESSES

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina



ON THE UPPER AND LOWER CLASSES FOR STATIONARY  
GAUSSIAN RANDOM FIELDS ON ABELIAN GROUPS WITH A REGULARLY VARYING ENTROPY

by

J.M.P. ALBIN

Technical Report No. 299

June 1990

000000 002

# ON THE UPPER AND LOWER CLASSES FOR STATIONARY GAUSSIAN RANDOM FIELDS ON ABELIAN GROUPS WITH A REGULARLY VARYING ENTROPY<sup>1</sup>

BY J. M. P. ALBIN

*Center for Stochastic Processes and University of Lund*

We give a complete and relatively explicit characterization of the upper and lower classes for a general stationary Gaussian random field.

**1. Introduction.** We shall assume that our probability space  $(\Omega, \mathcal{F}, \mathbf{P})$  is complete and that  $\{\xi(t)\}_{t \in T}$  is an  $\mathbb{R}$ -valued separable stochastically continuous standardized Gaussian random field on a pseudo-metric unbounded space  $(T, \rho)$  equipped with an abelian group-operation  $+$  such that the covariance  $r(s, t) \equiv \mathbf{E}\{\xi(s)\xi(t)\}$  satisfies  $r(s+u, t+u) = r(s, t)$  for  $s, t, u \in T$  and whose bounded subsets are totally bounded in the canonical pseudo-metric  $d(s, t) \equiv [\mathbf{E}\{(\xi(t) - \xi(s))^2\}]^{1/2}$ . We also define the entropy  $N_S(\varepsilon)$  as the minimum number of closed  $d$ -balls  $\mathcal{O}_\varepsilon$  of radius  $\varepsilon$  needed to cover  $S \subseteq T$  and  $M_S(\varepsilon)$  as the largest  $n$  for which there exist  $t_1, \dots, t_n \in S$  satisfying  $d(t_i, t_j) > \varepsilon$  for each  $i \neq j$ , and we write  $\mathbf{P}_0\{S\} \equiv \sup\{\mathbf{P}\{B\} : S \supseteq B \in \mathcal{F}\}$ ,  $\mathbf{P}^\circ\{S\} \equiv \inf\{\mathbf{P}\{B\} : S \subseteq B \in \mathcal{F}\}$ ,  $\Phi$  for the standard Gaussian d.f.,  $\underline{\Phi} \equiv 1 - \Phi$ ,  $0 \cdot \infty \equiv 0$ ,  $S_\rho(t, \varepsilon) \equiv \{s \in T : \rho(s, t) < \varepsilon\}$ ,  $S(t, \varepsilon) \equiv \{s \in T : d(s, t) \leq \varepsilon\}$  and  $\sigma(t, \varepsilon) \equiv \sup\{0 \vee r(s, t) : s \in T - S_\rho(t, \varepsilon)\}$ .

In view of recent tight tail-estimates for local suprema (over  $d$ -compact sets) of general Gaussian random fields (cf. e.g., [1], [2], [3], [16] and [21]), it seems motivated to study also the global behaviour of suprema. Here the only tractable approach seems to be upper and lower classes:

Let  $\Psi$  be the class of functions  $\psi : T \rightarrow [-\infty, \infty]$ . Provided that  $\sigma(t, \Delta) \rightarrow 0$  not too slowly as  $\Delta \rightarrow \infty$  we prove a zero-one law for the sets

$$E(\psi) \equiv \{\omega \in \Omega : \text{the set } \{t \in T : \xi(\omega; t) > \psi(t)\} \text{ is } \rho\text{-unbounded}\}, \psi \in \Psi.$$

We also give an explicit characterization of when the different values for  $\mathbf{P}\{E(\psi)\}$  occur, i.e., we determine the upper and lower classes for  $\xi(t)$ .

Related work are e.g., [5], [6], [10], [13], [14], [15], [17], [19] and [20].

**2. Main result.** Our main result is the following theorem.

**THEOREM 1.** Assume that there exists an  $R \in (0, \sqrt{2})$  such that

$$(2.1) \quad \overline{\lim}_{\varepsilon \downarrow 0} N_{\mathcal{O}_R}(x\varepsilon)/N_{\mathcal{O}_R}(\varepsilon) < \infty \quad \text{for some } x \in (0, 1),$$

<sup>1</sup>Supported by Air Force Office of Scientific Research Contract No. F49620 85C 0144 and by Kungliga Fysiografiska Sällskapet.

AMS 1980 subject classifications. Primary 60F15, 60F20, 60G10, 60G15, 60G17.

Key words and phrases. Upper and lower classes, LIL, Gaussian random fields.

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail. and/or Special
A-1	

and such that to each  $C > 0$  and  $s \in T$  there exists an increasing sequence  $\{\varrho_s(n)\}_{n=0}^\infty$ , with  $\varrho_s(0) = 0$  and  $\lim_{n \rightarrow \infty} \varrho_s(n) = \infty$  for  $s \in T$ , satisfying

$$(2.2) \sup_{s \in T} \sum_{\{n \geq 0 : \sigma(s, \varrho_s(n)) > 0\}} N_{S_\rho(s, \varrho_s(n+1))}(R) \exp\{-C/\sigma(s, \varrho_s(n))\} < \infty.$$

Then  $E(\psi) \in \mathcal{F}$  with  $\mathbf{P}\{E(\psi)\}$  zero or one for each  $\psi \in \Psi$  and moreover

$$(2.3) \mathbf{P}\{E(\psi)\} = 0 \Leftrightarrow \sum_{n=1}^\infty N_{\mathcal{O}_{r_n}}\left(\left(1 \vee \inf_{t \in S_n} \psi(t)\right)^{-1}\right) \Phi\left(1 \vee \inf_{t \in S_n} \psi(t)\right) < \infty$$

for some covering  $S_n = S(t_n, r_n)$ ,  $n = 1, 2, \dots$ , of  $T$  with  $r_n \leq R$  for all  $n$ .

REMARK 1. Note that, by (2.2), given  $\varepsilon > 0$  and  $t_0 \in T$ ,  $r(t, t_0) < \varepsilon$  for  $\rho(t, t_0) \geq k$  and  $k$  large, which yields  $S(t_0, \sqrt{2(1-\varepsilon)}) \subseteq S_\rho(t_0, k)$ . Thus  $\mathcal{O}_\delta$  is  $d$ -totally bounded for  $\delta < \sqrt{2}$  so that (2.1) makes sense and each covering  $\{S(t_n, r_n)\}$  of  $T$  with  $r_n \leq R$  is infinite.

PROOF:  $\Leftarrow$  We have, for  $\varepsilon \leq \delta \leq R/3$ , (since  $N_S(\varepsilon) \leq M_S(\varepsilon) \leq N_S(\varepsilon/2)$ ),

$$M_{\mathcal{O}_\delta}(\varepsilon) \leq N_{\mathcal{O}_\delta}(\varepsilon/2) \leq \frac{N_{\mathcal{O}_{R/3+\delta+\varepsilon}}(\varepsilon/2)}{M_{\mathcal{O}_{R/3}}(2\delta+2\varepsilon)} \leq \frac{N_{\mathcal{O}_R}(\varepsilon/2)}{N_{\mathcal{O}_R}(4\delta)/N_{\mathcal{O}_R}(R/3)},$$

and this inequality trivially extends to  $\varepsilon \leq \delta \leq R$ . Writing  $l$  for the smallest integer having  $x^{-l} \geq 8\delta/\varepsilon$  and  $K_1 \equiv \sup_{\varepsilon > 0} N_{\mathcal{O}_R}(x\varepsilon)/N_{\mathcal{O}_R}(\varepsilon)$  ( $< \infty$  by (2.1)), we readily get  $K_1^l \leq K_1(8\delta/\varepsilon)^{-\log K_1/\log x}$ , and thus

$$(2.4) \begin{aligned} M_{\mathcal{O}_\delta}(\varepsilon) &\leq N_{\mathcal{O}_\delta}(\varepsilon/2) \leq N_{\mathcal{O}_R}(R/3) \prod_{k=0}^{l-1} N_{\mathcal{O}_R}(4\delta x^{k+1})/N_{\mathcal{O}_R}(4\delta x^k) \\ &\leq K_1 N_{\mathcal{O}_R}(R/3) (8\delta/\varepsilon)^{-\log K_1/\log x} \text{ for } \varepsilon \leq \delta \leq R. \end{aligned}$$

Now, by (2.4),  $\overline{\lim}_{\varepsilon \downarrow 0} \log \log N_{\mathcal{O}_R}(\varepsilon)/\log(1/\varepsilon) = 0$  so  $\{\xi(t)\}_{t \in \mathcal{O}_R}$  has an a.s. bounded version; cf. [7], [8] and [18]. Since  $N_{S_\rho(t_0, \delta)}(R) < \infty$  for  $t_0 \in T$ ,  $\delta > 0$ ,  $\rho$ -separability yields that  $\{\xi(t)\}_{t \in S_\rho(t_0, \delta)}$  is a.s. bounded so

$$\mathbf{E}\{\sup_{t \in S_\rho(t_0, \delta)} \xi(t)^2\} \leq 2\mathbf{E}\{(\sup_{t \in S_\rho(t_0, \delta)} \xi(t))^2\} < \infty;$$

cf. [8], [9] and [11]. Since  $\xi(t)$  is stochastically continuous we get

$$d(t, t_0)^2 \leq \varepsilon^2 + \int_{G_\varepsilon} (\xi(t) - \xi(t_0))^2 d\mathbf{P} \leq \varepsilon^2 + 4 \int_{G_\varepsilon} \sup_{t \in S_\rho(t_0, \delta)} \xi(t)^2 d\mathbf{P} \rightarrow \varepsilon^2$$

as  $\rho(t, t_0) \rightarrow 0$ , where  $G_\varepsilon \equiv \{\omega \in \Omega : |\xi(\omega; t) - \xi(\omega; t_0)| > \varepsilon\}$ , so  $d(t, t_0) \rightarrow 0$  as  $\rho(t, t_0) \rightarrow 0$ . Hence  $d$ -opens are  $\rho$ -open and so  $\{\xi(t)\}_{t \in T}$  is  $d$ -separable. In view of  $\xi(t)$ 's (trivial)  $d$ -stochastic continuity it follows readily that any countable  $d$ -dense subset of  $\mathcal{O}_\varepsilon$  is a separator for  $\{\xi(t)\}_{t \in \mathcal{O}_\varepsilon}$ .

Take  $a_0 = \min\{(1-x^{1/2})^{1/2}/4, R/2\}$  and  $t \in T$ , let  $C_0 = \{t\}$  and let  $C_n$  be a  $(a/u)x^n$ -net in  $S(t, a/u)$  with  $d(s_1, s_2) > (a/u)x^n$  for  $C_n \ni s_1 \neq s_2 \in C_n$ , so  $\#C_n \leq M_{\mathcal{O}_{a/u}}((a/u)x^n)$ . Write  $p_n = (1-x^{1/2})x^{(n-1)/2}$  and  $C = \bigcup_{n=0}^\infty C_n$  and choose  $t_n(s) \in C_n$  with  $d(t_n(s), s) \leq (a/u)x^n$  for  $s \in C$ .

Then  $\xi(s) = \xi(t) + \sum_{n=1}^N [\xi(t_n(s)) - \xi(t_{n-1}(s))]$  for some  $N$  for each  $s \in C$ . Adapting [4, the proof of Theorem 6] to the present context we get

$$\begin{aligned} & \{\xi(s) > u + 1/u, \xi(t) \leq u\} \\ & \subseteq \bigcup_{n=1}^N \{\xi(t_n(s)) - \xi(t_{n-1}(s)) > p_n/u, \xi(t_n(s)) > u, \xi(t_{n-1}(s)) \leq u + 1/u\}. \end{aligned}$$

Thus, since  $d(t_n(s), t_{n-1}(s)) \leq d(t_n(s), s) + d(s, t_{n-1}(s)) \leq 2(a/u)x^{(n-1)}$ ,

$$\begin{aligned} (2.5) \quad & \mathbf{P}\{\sup_{s \in S(t, a/u)} \xi(s) > u + 1/u, \xi(t) \leq u\} \\ & = \mathbf{P}\{\bigcup_{s \in C} \{\xi(s) > u + 1/u\}, \xi(t) \leq u\} \\ & \leq \sum_{n=1}^{\infty} \sum_{s_1 \in C_{n-1}} \sum_{s_2 \in C_n \cap S(s_1, 2(a/u)x^{n-1})} \mathbf{P}\{\xi(s_2) - \xi(s_1) > p_n/u, \xi(s_2) > u, \xi(s_1) \leq u + 1/u\}. \end{aligned}$$

Now take  $a \in (0, a_0]$  and  $u \geq 1$  so that  $r(s_1, s_2) = 1 - d(s_1, s_2)^2/2 \geq 1 - 2(a/u)^2 \geq 1/2$  for  $d(s_1, s_2) \leq 2(a/u)x^{n-1}$ , which yields

$$\left(\frac{1}{r(s_1, s_2)} - 1\right)\xi(s_1) = \frac{d(s_1, s_2)^2}{2r(s_1, s_2)}\xi(s_1) \leq 4(a/u)^2 x^{2(n-1)} 2u \leq p_n/(2u)$$

for  $\xi(s_1) \leq u + 1/u$ . Hence we have, for  $d(s_1, s_2) \leq 2(a/u)x^{n-1}$ ,

$$\begin{aligned} (2.6) \quad & \mathbf{P}\{\xi(s_2) - \xi(s_1) > p_n/u, \xi(s_2) \geq u, \xi(s_1) \leq u + 1/u\} \\ & \leq \mathbf{P}\{\xi(s_2) - r(s_1, s_2)^{-1}\xi(s_1) > p_n/(2u), \xi(s_2) \geq u\} \\ & = \underline{\Phi}\left(\frac{\sqrt{2}r(s_1, s_2)p_n/(2u)}{\sqrt{1 + r(s_1, s_2)d(s_1, s_2)}}\right)\underline{\Phi}(u) \leq \underline{\Phi}\left(\frac{1 - x^{1/2}}{8ax^{(n-1)/2}}\right)\underline{\Phi}(u). \end{aligned}$$

Combining (2.4)-(2.6) we conclude that, uniformly for  $u \geq 1$ , as  $a \downarrow 0$ ,

$$\begin{aligned} (2.7) \quad & \underline{\Phi}(u)^{-1} \mathbf{P}\{\sup_{s \in S(t, a/u)} \xi(s) > u + 1/u, \xi(t) \leq u\} \\ & \leq \sum_{n=1}^{\infty} M_{\mathcal{O}_{a/u}}((a/u)x^{n-1}) M_{\mathcal{O}_{2(a/u)x^{n-1}}}((a/u)x^n) \underline{\Phi}\left(\frac{1 - x^{1/2}}{8ax^{(n-1)/2}}\right) \\ & \leq K_1^2 N_{\mathcal{O}_R}(R/3)^2 \sum_{n=1}^{\infty} (128x^{-n})^{-\log K_1 / \log x} \underline{\Phi}\left(\frac{1 - x^{1/2}}{8ax^{(n-1)/2}}\right) = o(a). \end{aligned}$$

Arguing as for (2.5) for  $\eta_u(s) \equiv 2u + 1/u - \xi(s)$  we deduce for future use that, by (2.4), (2.6) and symmetry, uniformly for  $u \geq 1$ , as  $a \downarrow 0$ ,

$$\begin{aligned} (2.8) \quad & \underline{\Phi}(u)^{-1} \mathbf{P}\{\inf_{s \in S(t, a/u)} \xi(s) < u, \xi(t) \geq u + 1/u\} \\ & = \underline{\Phi}(u)^{-1} \mathbf{P}\{\sup_{s \in S(t, a/u)} \eta_u(s) > u + 1/u, \eta_u(t) \leq u\} \end{aligned}$$

$$\begin{aligned}
&\leq \underline{\Phi}(u)^{-1} \sum_{n=1}^{\infty} \sum_{s_1 \in C_{n-1}} \sum_{s_2 \in C_n \cap S(s_1, 2(a/u)x^{n-1})} \\
&\quad \mathbf{P}\{\eta_u(s_2) - \eta_u(s_1) > p_n/u, \eta_u(s_2) > u, \eta_u(s_1) \leq u+1/u\} \\
&= \underline{\Phi}(u)^{-1} \sum_{n=1}^{\infty} \sum_{s_1 \in C_{n-1}} \sum_{s_2 \in C_n \cap S(s_1, 2(a/u)x^{n-1})} \\
&\quad \mathbf{P}\{\xi(s_1) - \xi(s_2) > p_n/u, \xi(s_1) \geq u, \xi(s_2) < u+1/u\} = o(a).
\end{aligned}$$

In order to proceed we observe that, by (2.4), for  $a \leq 1$  and  $\delta \leq R$ ,

$$\begin{cases} N_{\mathcal{O}_\delta}(a\varepsilon)/N_{\mathcal{O}_\delta}(\varepsilon) \leq N_{\mathcal{O}_\delta}(a\varepsilon) \leq K_1 N_{\mathcal{O}_R}(R/3)(8/a)^{-\log K_1/\log x}, \varepsilon \leq R, \\ N_{\mathcal{O}_\delta}(a\varepsilon) \leq N_{\mathcal{O}_R}(aR) \leq K_1 N_{\mathcal{O}_R}(R/3)(8/a)^{-\log K_1/\log x} N_{\mathcal{O}_\delta}(\varepsilon), \varepsilon > R. \end{cases}$$

Further  $u - 2/u \equiv \tilde{u} \geq \frac{1}{2}u \geq 1$  for  $u \geq 2$ , so that  $\tilde{u} + 1/\tilde{u} \leq u$ , and  $\underline{\Phi}(\tilde{u}) \leq \frac{1}{\tilde{u}}\phi(\tilde{u}) \leq \frac{2}{u}e^2\phi(u) \leq \frac{8}{3}e^2\underline{\Phi}(u)$ , where  $\phi(u) = (2\pi)^{-1/2} \exp\{-u^2/2\}$ . Now

$$\mathbf{P}\{\sup_{s \in S(t, a/u)} \xi(s) > u+1/u, \xi(t) \leq u\} \leq \underline{\Phi}(u) \quad \text{for } u \geq 1$$

for some sufficiently small  $a \in (0, a_0]$  (cf. (2.7)). Hence we conclude

$$\begin{aligned}
\mathbf{P}\left\{\sup_{s \in \mathcal{O}_\delta} \xi(s) > u\right\} &\leq N_{\mathcal{O}_\delta}(a/u) \left[ \mathbf{P}\left\{\sup_{s \in S(t, a/u)} \xi(s) > u, \xi(t) \leq \tilde{u}\right\} + \mathbf{P}\{\xi(t) > \tilde{u}\} \right] \\
&\leq N_{\mathcal{O}_\delta}(a/u) \left[ \mathbf{P}\left\{\sup_{s \in S(t, a/\tilde{u})} \xi(s) > \tilde{u} + 1/\tilde{u}, \xi(t) \leq \tilde{u}\right\} + \underline{\Phi}(\tilde{u}) \right] \\
&\leq \frac{16}{3}e^2 K_1 N_{\mathcal{O}_R}(R/3)(8/a)^{-\log K_1/\log x} N_{\mathcal{O}_\delta}(1/u) \underline{\Phi}(u)
\end{aligned}$$

for  $u \geq 2$ ,  $\delta \leq R$ , so, with  $K_2 = \frac{16}{3}e^2 K_1 N_{\mathcal{O}_R}(R/3)(8/a)^{-\log K_1/\log x}/\underline{\Phi}(2)$ ,

$$(2.9) \quad \mathbf{P}\left\{\sup_{s \in \mathcal{O}_\delta} \xi(s) > u\right\} \leq K_2 N_{\mathcal{O}_\delta}(1/(1 \vee u)) \underline{\Phi}(1 \vee u) \quad \text{for } \delta \leq R \text{ and all } u.$$

Assume that the sum (2.3) is finite for a covering  $\{S_n\} = \{S(t_n, r_n)\}$  of  $T$  with  $r_n \leq R$ . Taking  $m = \sup\{\rho(t_1, t_n) : 1 \leq n < J\}$  where

$$\sum_{n=J}^{\infty} N_{\mathcal{O}_{r_n}}((1 \vee \inf_{t \in S_n} \psi(t))^{-1}) \underline{\Phi}(1 \vee \inf_{t \in S_n} \psi(t)) < \varepsilon/K_2,$$

completeness yields that  $E(\psi) \in \mathcal{F}$  with  $\mathbf{P}\{E(\psi)\} = 0$  since, by (2.9),

$$\begin{aligned}
\mathbf{P}^\circ\{E(\psi)\} &\leq \mathbf{P}^\circ\{\xi(t) > \psi(t), \text{ for some } t \in T \text{ with } \rho(t_1, t) > m+R\} \\
&\leq \mathbf{P}\left\{\bigcup_{\{n: \rho(t_1, t_n) > m\}} \{\xi(t) > \inf_{s \in S_n} \psi(s), \text{ for some } t \in S_n\}\right\} \\
&\leq K_2 \sum_{n=J}^{\infty} N_{\mathcal{O}_{r_n}}\left((1 \vee \inf_{t \in S_n} \psi(t))^{-1}\right) \underline{\Phi}\left(1 \vee \inf_{t \in S_n} \psi(t)\right) < \varepsilon.
\end{aligned}$$

$\Rightarrow$  Write  $\sum(\{S_n\}; \psi)$  for the sum (2.3) and assume that  $\sum(\{S_n\}; \psi) = \infty$  for each covering  $S_n = S(t_n, r_n)$ ,  $n = 1, 2, \dots$ , of  $T$  with  $r_n \leq R$ .

Taking  $t_0 \in T$  and  $2 \leq u_1 \leq u_2 \leq \dots$  with  $\mathbf{P}\{\sup_{t \in S_\rho(t_0, n)} \xi(t) > u_n\} \leq n^{-2}$  (recall that  $\{\xi(t)\}_{t \in S_\rho(t_0, n)}$  is a.s. bounded), the function  $\psi^*(t) \equiv u_1$  for  $t \in S_\rho(t_0, 1)$  and  $\psi^*(t) \equiv u_n$  for  $t \in S_\rho(t_0, n) - S_\rho(t_0, n-1)$ ,  $n \geq 2$ , has

$$\mathbf{P}^\circ\{E(\psi^*)\} \leq \lim_{n \rightarrow \infty} \mathbf{P}^\circ\{\xi(t) > \psi^*(t), \text{ for some } t \in T - S_\rho(t_0, n)\} = 0.$$

Clearly  $\mathbf{P}_\circ\{A \cup B\} \leq \mathbf{P}^\circ\{A\} + \mathbf{P}_\circ\{B\}$  so that  $\mathbf{P}_\circ\{E(\psi \wedge \psi^*)\} = \mathbf{P}_\circ\{E(\psi) \cup E(\psi^*)\} \leq \mathbf{P}_\circ\{E(\psi)\}$  and so, by completeness, it suffices to prove that

$$(2.10) \quad \varphi(t) \equiv (\psi(t) \wedge \psi^*(t)) \vee 2 \quad \text{has} \quad \mathbf{P}_\circ\{E(\varphi)\} = 1.$$

Take a  $(p/u)$ -net  $\{s_i\}_{i=1}^n$  in  $\mathcal{O}_\delta$  with  $d(s_i, s_j) > p/u$  for  $s_i \neq s_j$ . Since

$$(2.11) \quad \begin{aligned} M_{\mathcal{O}_{\delta \wedge (kp/u)}}(p/u) &= M_{\mathcal{O}_{\delta \wedge (kp/u)}}((2\delta) \wedge (p/u)) \\ &\leq K_1 N_{\mathcal{O}_R}(R/3) \left(8 \frac{\delta \wedge (kp/u)}{(2\delta) \wedge (p/u)}\right)^{-\log K_1 / \log x} \\ &\leq K_1 N_{\mathcal{O}_R}(R/3) (8k)^{-\log K_1 / \log x} \quad \text{for } \delta \leq R, k \geq 1 \end{aligned}$$

(again using (2.4)), and since, by arguing as for [5, Eq. 2.16], for  $x, y > 0$ ,

$$(2.12) \quad \mathbf{P}\{\xi(s) > x, \xi(t) > y\} \leq \underline{\Phi}(\tfrac{1}{2}d(s, t)x) \underline{\Phi}(y) + \underline{\Phi}(\tfrac{1}{2}d(s, t)y) \underline{\Phi}(x)$$

for all values of  $r(s, t)$  (although [5] only treat  $0 \leq r(s, t) < 1$ ), we obtain

$$\begin{aligned} &\sum_{i \neq j} \mathbf{P}\{\xi(s_i) > u, \xi(s_j) > u\} \\ &\leq 2 \underline{\Phi}(u) \sum_{i=1}^n \sum_{k=1}^{\lfloor 2\delta u/p \rfloor} \sum_{\{1 \leq j \leq n: kp/u < d(s_i, s_j) \leq (k+1)p/u\}} \underline{\Phi}(\tfrac{1}{2}d(s_i, s_j)u) \\ &\leq 2n \underline{\Phi}(u) K_1 N_{\mathcal{O}_R}(R/3) \sum_{k=1}^{\infty} (8(k+1))^{-\log K_1 / \log x} \underline{\Phi}(\tfrac{1}{2}kp) \leq \tfrac{1}{2}n \underline{\Phi}(u) \end{aligned}$$

for  $u > 0$ ,  $\delta \leq R$  and for some  $p \geq 1$  (not depending on  $\delta$ ). Since, by (2.4),

$$\begin{aligned} N_{\mathcal{O}_\delta}(1/u) &\leq N_{\mathcal{O}_{\delta \wedge (p/u)}}(\delta \wedge (1/u)) N_{\mathcal{O}_\delta}(\delta \wedge (p/u)) \\ &\leq K_1 N_{\mathcal{O}_R}(R/3) (8p)^{-\log K_1 / \log x} n \quad \text{for } \delta \leq R \end{aligned}$$

we conclude, taking  $K_3 = \tfrac{1}{2} K_1^{-1} N_{\mathcal{O}_R}(R/3)^{-1} (8p)^{\log K_1 / \log x}$ ,

$$(2.13) \quad \begin{aligned} \mathbf{P}\{\sup_{t \in \mathcal{O}_\delta} \xi(t) > u\} &\geq \mathbf{P}\{\sup_{1 \leq i \leq n} \xi(s_i) > u\} \\ &\geq n \underline{\Phi}(u) - \sum_{i \neq j} \mathbf{P}\{\xi(s_i) > u, \xi(s_j) > u\} \\ &\geq K_3 N_{\mathcal{O}_\delta}(1/u) \underline{\Phi}(u) \quad \text{for } u > 0 \text{ and } \delta \leq R. \end{aligned}$$

Now, combining (2.9) and (2.13) we get, for each choice of  $\{S_n\}$ ,

$$(2.14) \quad K_2 \sum(\{S_n\}; \varphi) \geq \sum_{n=1}^{\infty} \mathbf{P}\left\{\sup_{t \in S_n} \xi(t) > \inf_{t \in S_n} \varphi(t)\right\}$$

$$\begin{aligned} &\geq \sum_{n=1}^{\infty} \mathbf{P}\left\{\sup_{t \in S_n} \xi(t) > \inf_{t \in S_n} \psi(t) \wedge \psi^*(t)\right\} \mathbf{P}\left\{\sup_{t \in S_n} \xi(t) > 2\right\} \\ &\geq K_3 \underline{\Phi}(2) \sum(\{S_n\}; \psi) = \infty. \end{aligned}$$

Let  $r_t \equiv \sup\{r > 0 : r \inf_{s \in S(t,r)} \varphi(s) < a\}$  for  $a \in (0, 1]$ ,  $t \in T$ , so that  $a/\psi^*(t) \leq r_t \leq a/2$ . Taking  $\delta_k \uparrow r_t$  with  $\delta_k \inf_{s \in S(t, \delta_k)} \varphi(s) < a$  we get

$$(2.15) \quad \begin{cases} a/(\inf_{s \in S(t, r_t)} \varphi(s)) \geq \lim_{k \rightarrow \infty} a/(\inf_{s \in S(t, \delta_k)} \varphi(s)) \geq \lim_{k \rightarrow \infty} \delta_k = r_t, \\ a/(\inf_{s \in S(t, r_t)} \varphi(s)) \leq \lim_{\varepsilon \downarrow 0} a/(\inf_{s \in S(t, r_t + \varepsilon)} \varphi(s)) \leq \lim_{\varepsilon \downarrow 0} r_t + \varepsilon = r_t. \end{cases}$$

Ordering  $\mathcal{S} \equiv \{A \subseteq T : A \ni s \neq t \in A \Rightarrow d(s, t) > r_s \wedge r_t\}$  partially by  $A \leq B \Leftrightarrow A \subseteq B$ , a chain  $\{A_\alpha\} \subseteq \mathcal{S}$  has upper bound  $\cup\{A_\alpha\}$  so that, by Zorn's Lemma,  $\mathcal{S}$  has a maximal element  $\mathcal{C}$ . Here  $\mathcal{C}$ 's maximality readily yields  $\cup_{t \in \mathcal{C}} S_t = T$ , where  $S_t \equiv S(t, r_t)$ . Further, since  $\#\mathcal{C} \cap S_\rho(t_0, n) \leq M_{S_\rho(t_0, n)}(a/u_n) < \infty$ , we have  $\#\mathcal{C} \leq \aleph_0$  and, by (2.14),  $\sum(\{S_t\}; \psi) = \infty$ . Writing  $\varphi_t = \inf_{s \in S_t} \varphi(s)$  we therefore obtain, by (2.4) and (2.15),

$$(2.16) \quad \sum_{t \in \mathcal{C}} \underline{\Phi}(\varphi_t) \geq \frac{(8/a) \log K_1 / \log x}{K_1 N_{\mathcal{C}_R}(R/3)} \sum_{t \in \mathcal{C}} N_{S_t}(ar_t) \underline{\Phi}(\varphi_t) = \infty.$$

Now let  $\varphi_t^* \equiv \varphi_t + 1/\varphi_t$ ,  $J_t \equiv \{\omega \in \Omega : \xi(\omega; t) > \varphi_t^*, \inf_{s \in S_t} \xi(\omega; s) \geq \varphi_t\}$  and  $\mathcal{C}_m^N \equiv \{t \in \mathcal{C} : m \leq \rho(t_0, t) < N\}$ . Letting  $I_t$  "indicate"  $J_t$  we get

$$\begin{aligned} (2.17) \quad \mathbf{P}_0\{E(\varphi)\} &= \mathbf{P}_0\left\{\bigcap_{m=1}^{\infty} \bigcup_{N=m}^{\infty} \bigcup_{t \in \mathcal{C}_m^N} \{\xi(\omega; s) > \varphi(s), \text{ for some } s \in S_t\}\right\} \\ &\geq \mathbf{P}\left\{\bigcap_{m=1}^{\infty} \bigcup_{N=m}^{\infty} \left\{\sum_{t \in \mathcal{C}_m^N} I_t > 0\right\}\right\} \\ &\geq \overline{\lim}_{m \rightarrow \infty} \overline{\lim}_{N \rightarrow \infty} \left[1 + \text{Var}\left\{\sum_{t \in \mathcal{C}_m^N} I_t\right\} / \left(\mathbf{E}\left\{\sum_{t \in \mathcal{C}_m^N} I_t\right\}\right)^2\right]^{-1}, \end{aligned}$$

where we used Hölder's inequality as in [5]. Now write

$\mu_{s,t} = \mathbf{P}\{\xi(s) > \varphi_s^*, \xi(t) > \varphi_t^*\} - \mathbf{P}\{\xi(s) > \varphi_s^*\} \mathbf{P}\{\xi(t) > \varphi_t^*\}$  for  $s, t \in \mathcal{C}$  and note that, by arguing as above,  $\underline{\Phi}(\varphi_t^*) \geq \frac{3}{8} e^{-2} \underline{\Phi}(\varphi_t)$  so that, by (2.8),

$\mathbf{E}\{I_t\} = \underline{\Phi}(\varphi_t^*) - \mathbf{P}\{\xi(t) > \varphi_t^*, \inf_{s \in S_t} \xi(s) < \varphi_t\} \geq \frac{3}{16} e^{-2} \underline{\Phi}(\varphi_t)$  for  $t \in \mathcal{C}$  and  $a \leq a_1$ , for some  $a_1 \in (0, R]$  (not depending on  $t$ ). Since, by (2.8),

$$\begin{aligned} \text{Var}\left\{\sum_{t \in \mathcal{C}_m^N} I_t\right\} &\leq \sum_{s, t \in \mathcal{C}_m^N} \left[\mu_{s,t} + 2 \underline{\Phi}(\varphi_s^*) \mathbf{P}\{\xi(t) > \varphi_t^*, \inf_{v \in S_t} \xi(v) < \varphi_t\}\right] \\ &= \sum_{s, t \in \mathcal{C}_m^N} \mu_{s,t} + o(a) \left(\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t^*)\right)^2 \end{aligned}$$

(see also [5]), (2.16) and (2.17) combines to show that it suffices to prove



$$(2.18) \quad \lim_{m \rightarrow \infty} \lim_{N \rightarrow \infty} (\sum_{s,t \in C_m^N} \mu_{s,t}) / (\sum_{t \in C_m^N} \Phi(\varphi_t))^2 \leq 0 \text{ for } a \leq a_1.$$

Now, given an integer  $k \geq 1$ , partition  $C_m^N \times C_m^N$  into

$$\begin{cases} C_{m,N}^{k,1} \equiv \{(s,t) : d(s,t) > R, 0 < r(s,t) \leq k^{-1}[(\varphi_s^*)^2 + (\varphi_t^*)^2]^{-1}\}, \\ C_{m,N}^{k,2} \equiv \{(s,t) : d(s,t) > R, r(s,t) > k^{-1}[(\varphi_s^*)^2 + (\varphi_t^*)^2]^{-1}\}, \\ C_{m,N}^3 \equiv \{(s,t) : 0 < d(s,t) \leq R, r(s,t) > 0, \frac{1}{2}\varphi_s \leq \varphi_t \leq 2\varphi_s\}, \\ C_{m,N}^4 \equiv \{(s,t) : 0 < d(s,t) \leq R, r(s,t) > 0, \varphi_t > 2\varphi_s \text{ or } \varphi_s > 2\varphi_t\}, \\ C_{m,N}^5 \equiv \{(s,t) : d(s,t) > 0, r(s,t) \leq 0\}, \\ C_{m,N}^6 \equiv \{(s,t) : s=t\}. \end{cases}$$

Arguing as for [5, Eq. 2.12] we then readily get

$$(2.19) \quad \mu_{s,t} \leq \frac{e^{1/(2k)} \phi(\varphi_s^*) \phi(\varphi_t^*)}{\sqrt{2Rk} \varphi_s^* \varphi_t^*} \leq \frac{16 e^{1/(2k)} \Phi(\varphi_s) \Phi(\varphi_t)}{9 \sqrt{2Rk}} \text{ for } (s,t) \in C_{m,N}^{k,1}.$$

Further we have, by arguing as for [5, Eq. 2.13],

$$\mu_{s,t} \leq \frac{\varphi_s^* \phi(\varphi_s^*)}{\sqrt{\pi R}} \exp\left\{-\frac{R^2(\varphi_t^*)^2}{8}\right\} \text{ for } \varphi_t^* \geq \varphi_s^* \text{ and } d(s,t) > R.$$

Observing that  $x \exp\{-Cx^2\} \leq (2C)^{-1/2}$  and taking  $\varrho$  corresponding to  $C \equiv R^2/(48k)$  in (2.2) we thus obtain, by (2.4) and (2.15), for  $s \in C_m^N$ ,

$$\begin{aligned} & \sum_{\{t \in C_m^N : (s,t) \in C_{m,N}^{k,2}, \varphi_t^* \geq \varphi_s^*\}} \mu_{s,t} \\ & \leq \frac{4 \Phi(\varphi_s)}{3 \sqrt{\pi R}} \sum_{\ell=2}^{\infty} \sum_{n=0}^{\infty} \sum_{\{t \in C_m^N : \ell \leq \varphi_t^* < \ell+1, \varrho_s(n) \leq \varrho(s,t) < \varrho_s(n+1), r(s,t) > 0\}} \\ & \quad \varphi_t^* \exp\left\{-\frac{R^2(\varphi_t^*)^2}{12}\right\} \exp\left\{-\frac{r^2}{48k r(s,t)}\right\} \\ & \leq \frac{8 \Phi(\varphi_s)}{\sqrt{3\pi R^2}} \sum_{\ell=2}^{\infty} \sum_{\{n \geq 0 : \sigma(s, \varrho_s(n)) \geq 0\}} M_{S_{\varrho}(s, \varrho_s(n+1))}(a/(\ell+1)) \\ & \quad \times \exp\left\{-\frac{R^2 \ell^2}{24}\right\} \exp\left\{-\frac{C}{\sigma(s, \varrho_s(n))}\right\} \\ & \leq \frac{8 \Phi(\varphi_s)}{\sqrt{3\pi R^2}} \sum_{\{n \geq 0 : \sigma(s, \varrho_s(n)) \geq 0\}} N_{S_{\varrho}(s, \varrho_s(n+1))}(R) \exp\left\{-\frac{C}{\sigma(s, \varrho_s(n))}\right\} \\ & \quad \times K_1 N_{\mathcal{O}_R}(R/3) \sum_{\ell=2}^{\infty} (8R(\ell+1)/a)^{-\log K_1 / \log x} \exp\left\{-\frac{R^2 \ell^2}{24}\right\}. \end{aligned}$$

Since  $\sum_{t \in \mathcal{C}_0^m} \underline{\Phi}(\varphi_t) \leq N_{S_p(t_0, m)}(a/u_m) \underline{\Phi}(u_m) < \infty$  so that, by (2.16),  $\lim_{N \rightarrow \infty} \sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t) = \infty$ , we readily deduce, by (2.2) and symmetry,

$$(2.20) \quad \lim_{N \rightarrow \infty} (\sum_{(s,t) \in \mathcal{C}_{m,N}^{k,2}} \mu_{s,t}) / (\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t))^2 = 0 \quad \text{for } a \leq a_1.$$

Clearly we have, by (2.12) and (2.15), for  $s \in \mathcal{C}_m^N$ ,

$$\begin{aligned} & \sum_{\{t \in \mathcal{C}_m^N : (s,t) \in \mathcal{C}_{m,N}^3, \varphi_t \geq \varphi_s\}} \mu_{s,t} \\ & \leq \sum_{\ell=1}^{\infty} \sum_{\{t \in \mathcal{C}_m^N : \ell a / (2\varphi_s) < d(s,t) \leq R \wedge ((\ell+1)a / (2\varphi_s)), \varphi_t \leq 2\varphi_s\}} \\ & \quad 2 \underline{\Phi}(\varphi_s) \underline{\Phi}(\tfrac{1}{2}d(s,t)\varphi_s) \\ & \leq 2 \underline{\Phi}(\varphi_s) \sum_{\ell=1}^{\infty} M_{\mathcal{O}_{R \wedge ((\ell+1)a/(2\varphi_s))}}(a/(2\varphi_s)) \underline{\Phi}(\tfrac{1}{4}\ell a), \end{aligned}$$

and using (2.11) and symmetry we thus get (since  $\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t) \rightarrow \infty$ )

$$(2.21) \quad \lim_{N \rightarrow \infty} (\sum_{(s,t) \in \mathcal{C}_{m,N}^3} \mu_{s,t}) / (\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t))^2 = 0 \quad \text{for } a \leq a_1.$$

Further we have, for  $s \in \mathcal{C}_m^N$ , by [12, Theorem 4.2.1] and using (2.4), (2.15) and the facts that  $\varphi_s \geq 2$  and that  $x^\beta \exp\{-Cx^2\} \leq (\beta/(2C))^{\beta/2}$ ,

$$\begin{aligned} & \sum_{\{t \in \mathcal{C}_m^N : (s,t) \in \mathcal{C}_{m,N}^4, \varphi_t > 2\varphi_s\}} \mu_{s,t} \\ & \leq \sum_{\ell=2}^{\infty} \sum_{\{t \in \mathcal{C}_m^N : \ell\varphi_s < \varphi_t \leq (\ell+1)\varphi_s, r(s,t) > 0, 0 < d(s,t) \leq R\}} \\ & \quad \frac{r(s,t)}{\sqrt{2\pi} d(s,t) \sqrt{1+r(s,t)}} \exp\left\{-\frac{(\varphi_s^*)^2 + (\varphi_t^*)^2}{2(1+r(s,t))}\right\} \\ & \leq \sum_{\ell=2}^{\infty} \frac{(\ell+1)\varphi_s}{\sqrt{2\pi} a} M_{\mathcal{O}_R}(a/((\ell+1)\varphi_s)) \exp\left\{-\frac{(\ell^2+1)\varphi_s^2}{4}\right\} \\ & \leq \underline{\Phi}(\varphi_s) \sum_{\ell=2}^{\infty} \frac{4K_1 N_{\mathcal{O}_R}(R/3)(\ell+1)\varphi_s^2}{3\sqrt{\pi} a (8R(\ell+1)\varphi_s/a)^{\log K_1 / \log x}} \exp\left\{-\frac{(\ell^2-1)\varphi_s^2}{4}\right\} \\ & \leq \underline{\Phi}(\varphi_s) \sum_{\ell=2}^{\infty} \frac{N_{\mathcal{O}_R}(R/3)(2 - \frac{\log K_1}{\log x})^{1 - \log K_1 / \log x}}{6\sqrt{\pi} R (8R(\ell+1)/a)^{\log K_1 / \log x - 1}} \exp\{-(\ell^2-3)\}, \end{aligned}$$

and using symmetry we thus get (again since  $\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t) \rightarrow \infty$ )

$$(2.22) \quad \lim_{N \rightarrow \infty} (\sum_{(s,t) \in \mathcal{C}_{m,N}^+} \mu_{s,t}) / (\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t))^2 = 0 \quad \text{for } a \leq a_1.$$

Finally, by [12, Theorem 4.2.1],  $\mu_{s,t} \leq 0$  for  $r(s,t) \leq 0$ ,  $s \neq t$ , so that

$$(2.23) \quad \lim_{N \rightarrow \infty} (\sum_{(s,t) \in \mathcal{C}_{m,N}^+} \mu_{s,t}) / (\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t))^2 \leq 0 \quad \text{for } a \leq a_1,$$

and moreover

$$(2.24) \quad \lim_{N \rightarrow \infty} \frac{\sum_{(s,t) \in \mathcal{C}_{m,N}^+} \mu_{s,t}}{(\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t))^2} = \lim_{N \rightarrow \infty} \frac{1}{\sum_{t \in \mathcal{C}_m^N} \underline{\Phi}(\varphi_t)} = 0 \quad \text{for } a \leq a_1.$$

Combining (2.19)-(2.24) we see that (given  $a < a_1$ ) the left hand side of (2.18) is at most  $\mathcal{O}(1/k)$ , and so (2.18) follows from sending  $k \uparrow \infty$ .

**COROLLARY 1.** Assume the hypothesis of Theorem 1 and that  $d$  is a metric and  $(T, d)$  locally compact. Then there exists an invariant (w.r.t.  $+$ ) Haar-measure  $\mu$  on  $(T, d)$ 's Borel-sets with  $\mu(\mathcal{O}_\delta) < \infty$  for  $\delta \in (0, \sqrt{2})$ . If further  $\lambda$  is any version of this Haar-measure, then  $\mathbf{P}\{E(\psi)\} = 0$  i.f.f.

$$(2.25) \quad \sum_{n=1}^{\infty} \left[ 1 + \lambda(\mathcal{O}_{r_n}) N_{\mathcal{O}_R} \left( \left( 1 \vee \inf_{t \in S_n} \psi(t) \right)^{-1} \right) \right] \underline{\Phi} \left( 1 \vee \inf_{t \in S_n} \psi(t) \right) < \infty$$

for some covering  $S_n = S(t_n, r_n)$ ,  $n = 1, 2, \dots$ , of  $T$  with  $r_n \leq R$  for all  $n$ .

**PROOF:** An easy argument yields  $d$ -continuity for  $(s, t) \rightarrow t - s$  so that  $(T, d, +)$  is a locally compact (Hausdorff) topological group and  $\mu$  exists and is finite on compacts, where, by Remark 1 and local compactness,  $\mathcal{O}_\delta$  is compact for  $\delta < \sqrt{2}$ . Now observe that, by arguing as for (2.4),

$$N_{\mathcal{O}_\delta}(\varepsilon) \leq 1 + \frac{K_1 N_{\mathcal{O}_R}(R/3)^2 N_{\mathcal{O}_R}(\varepsilon)}{32^{\log K_1 / \log x} N_{\mathcal{O}_R}(\delta)} \leq 1 + \frac{K_1 N_{\mathcal{O}_R}(R/3)^2 \lambda(\mathcal{O}_\delta) N_{\mathcal{O}_R}(\varepsilon)}{32^{\log K_1 / \log x} \lambda(\mathcal{O}_R)}$$

for  $\varepsilon > 0$  and  $\delta \leq R$ , and so the sum (2.3) is finite when (2.25) holds. Conversely (2.25) holds when the sum (2.3) is finite since, for  $\delta \leq R$ ,

$$\begin{aligned} \frac{N_{\mathcal{O}_R}(\varepsilon)}{N_{\mathcal{O}_\delta}(\varepsilon)} &\leq N_{\mathcal{O}_R}(R/2) M_{\mathcal{O}_{R/2}}((R/2) \wedge (2\delta)) N_{\mathcal{O}_{(R/2) \wedge (2\delta)}}(\delta) \\ &\leq \frac{K_1 N_{\mathcal{O}_R}(R/3) N_{\mathcal{O}_R}(R/2) \lambda(\mathcal{O}_R)}{16^{\log K_1 / \log x} \lambda(\mathcal{O}_{(R/4) \wedge \delta})} \leq \frac{K_1 N_{\mathcal{O}_R}(R/3)^2 \lambda(\mathcal{O}_R)^2}{16^{\log K_1 / \log x} \lambda(\mathcal{O}_{R/4}) \lambda(\mathcal{O}_\delta)}. \end{aligned}$$

The following local result improves on [16] and [21] (but note that they also treat non-stationary fields); we leave to the reader to find what conditions in Section 1 one can omit without violating it's conclusion.

**COROLLARY 2.** Assume that there exists an  $R \in (0, \sqrt{2})$  such that (2.1) holds. Then there exist constants  $C_1, C_2 \in (0, \infty)$  such that

$$C_1 \leq \frac{\mathbf{P}\{\sup_{t \in \mathcal{O}_\delta} \xi(t) > u\}}{N_{\mathcal{O}_\delta}((1 \vee u)^{-1}) \underline{\Phi}(u)} \leq C_2 \quad \text{for } u \in \mathbb{R} \text{ and } \delta \in [0, R].$$

If in addition  $d$  is a metric,  $(T, d)$  is locally compact and  $\lambda$  is a version of the Haar-measure, then there exist constants  $C_1, C_2 \in (0, \infty)$  such that

$$C_1 \leq \frac{\mathbf{P}\{\sup_{t \in \mathcal{O}_\delta} \xi(t) > u\}}{[1 + \lambda(\mathcal{O}_\delta) N_{\mathcal{O}_R}((1 \vee u)^{-1})] \underline{\Phi}(u)} \leq C_2 \quad \text{for } u \in \mathbb{R} \text{ and } \delta \in [0, R].$$

For homogeneous spaces which in a certain sense are finite-dimensional we have the following very simple criterion for (2.2) to hold.

**PROPOSITION 1.** *If  $\rho(s+u, t+u) = \rho(s, t)$  for  $s, t, u \in T$  and if there exists a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that, writing  $\mathcal{B}_\varepsilon$  for an open  $\rho$ -ball of radius  $\varepsilon$ ,*

$$(2.26) \quad 1 < \lim_{\Delta \rightarrow \infty} \frac{N_{\mathcal{B}_{\Delta+f(\Delta)}}(R)}{N_{\mathcal{B}_\Delta}(R)} \leq \overline{\lim}_{\Delta \rightarrow \infty} \frac{N_{\mathcal{B}_{\Delta+f(\Delta)}}(R)}{N_{\mathcal{B}_\Delta}(R)} < \infty,$$

then (2.2) holds if  $\sigma(\varepsilon) \equiv \sup\{0 \vee r(s, t) : \rho(s, t) \geq \varepsilon\}$  satisfies

$$(2.27) \quad \lim_{\Delta \rightarrow \infty} \sigma(\Delta) \log N_{\mathcal{B}_\Delta}(R) = 0.$$

**PROOF:** Take  $\varepsilon, y, \Delta > 0$  with  $1 + \varepsilon \leq N_{\mathcal{B}_{x+f(x)}}(R)/N_{\mathcal{B}_x}(R) \leq y$  for  $x \geq \Delta$  and let  $\varrho(0) = 0$ ,  $\varrho(1) = \Delta$  and  $\varrho(n+1) = \varrho(n) + f(\varrho(n))$  for  $n \geq 1$ , so that

$$N_{\mathcal{B}_{\varrho(n+1)}}(R)/N_{\mathcal{B}_{\varrho(n)}}(R) = \prod_{k=1}^n N_{\mathcal{B}_{\varrho(k+1)}}(R)/N_{\mathcal{B}_{\varrho(k)}}(R) \geq (1 + \varepsilon)^n \rightarrow \infty$$

as  $n \rightarrow \infty$ , which yields  $\lim_{n \rightarrow \infty} \varrho(n) = \infty$ . Taking  $n_0$  such that  $\sigma(\varrho(n)) \times \log N_{\mathcal{B}_{\varrho(n)}}(R) \leq C/2$  for  $n \geq n_0$  we now readily obtain

$$\begin{aligned} \sup_{s \in T} \sum_{\{n \geq 0 : \sigma(s, \varrho(n)) > 0\}} N_{\mathcal{B}_{\varrho(n+1)}}(R) \exp\{-C/\sigma(s, \varrho(n))\} \\ \leq \sum_{n=1}^{n_0} N_{\mathcal{B}_{\varrho(n)}}(R) + \sum_{n=n_0}^{\infty} N_{\mathcal{B}_{\varrho(n+1)}}(R) \exp\{-2 \log N_{\mathcal{B}_{\varrho(n)}}(R)\} \\ \leq \sum_{n=1}^{n_0} N_{\mathcal{B}_{\varrho(n)}}(R) + \sum_{n=n_0}^{\infty} y N_{\mathcal{B}_{\varrho(n)}}(R)^{-1} (1 + \varepsilon)^{-(n-1)} < \infty. \end{aligned}$$

**3. The Euclidian case.** Theorem 2 sharpens [10] and [15]: They need a stronger condition than (3.1) and only treat (and crucially need)  $\psi(t) = \hat{\psi}(|t|)$  with  $\hat{\psi}: [0, \infty) \rightarrow [0, \infty)$  increasing (a meaningless notion on general space), which makes (3.3) an integraltest and proofs totally different.

**THEOREM 2.** *If  $\{\xi(t)\}_{t \in \mathbb{R}^n}$  is separable stationary standard Gaussian, if*

$$(3.1) \quad \lim_{|t-s| \rightarrow \infty} (0 \vee r(s, t)) \log |t-s| = 0,$$

and if there exist constants  $\alpha, \delta, C_1, C_2 \in (0, \infty)$  such that

$$(3.2) \quad C_1 |t-s|^\alpha \leq 1 - r(s, t) \leq C_2 |t-s|^\alpha \quad \text{for } 0 \leq |t-s| \leq \delta,$$

then  $E(\psi) \in \mathcal{F}$  with  $\mathbf{P}\{E(\psi)\}$  zero or one for each  $\psi \in \Psi$  and moreover, writing  $\lambda$  for the Lebesgue measure over  $\mathbb{R}^n$ , we have  $\mathbf{P}\{E(\psi)\} = 0$  i.f.f.

$$(3.3) \quad \sum_{n=1}^{\infty} \left[ 1 + \lambda(\mathcal{O}_{r_n}) \left( 1 \vee \inf_{t \in S_n} \psi(t) \right)^{2n/\alpha} \right] \underline{\Phi} \left( 1 \vee \inf_{t \in S_n} \psi(t) \right) < \infty$$

for some covering  $S_n = S(t_n, r_n)$ ,  $n = 1, 2, \dots$ , of  $T$  with  $r_n \leq 1$  for all  $n$ .

PROOF: Here  $(T, \rho, +) = (\mathbb{R}^n, |\cdot|, +)$  and  $R = 1$ . Take  $\Delta > 0$  with  $r(0, t) < \frac{1}{2}$  for  $|t| \geq \Delta$  and suppose  $|t| \not\rightarrow 0$  as  $d(0, t) \rightarrow 0$ . Then  $\inf\{d(0, t) : |t| \geq \varrho\} = 0$  for some  $\varrho \in (0, \Delta]$ , and picking  $s$  with  $|s| \geq \varrho$  and  $d(0, s) < \frac{\varrho}{2\Delta}$  we get  $d(0, ([\frac{\Delta}{\varrho}] + 1)s) < 1$  so  $r(0, ([\frac{\Delta}{\varrho}] + 1)s) > \frac{1}{2}$ . This is a contradiction since  $|\frac{\Delta}{\varrho}s| \geq \Delta$ , and so, by homogeneity,  $|t - s| \rightarrow 0$  as  $d(s, t) \rightarrow 0$ . Now pick  $\varrho > 0$  with  $|t - s| \leq \delta$  for  $d(s, t) \leq \varrho$ . Then (3.1) readily yields that

$$(3.4) \quad S_{|\cdot|}(t, (2C_2)^{-1/\alpha} \varepsilon^{2/\alpha}) \subseteq S(t, \varepsilon) \subseteq S_{|\cdot|}(t, C_1^{-1/\alpha} \varepsilon^{2/\alpha}) \quad \text{for } \varepsilon \leq \delta \wedge \varrho.$$

Thus  $|\cdot|$ - and  $d$ -topologies coincide so we have stochastic continuity,  $|\cdot|$ -boundeds are  $d$ -totally bounded,  $d$  is a metric,  $(T, d)$  is locally compact and the Lebesgue measure is a Haar-measure on  $(T, d, +)$ . Further it follows easily from (3.4), since  $S(t, 1) \subseteq S_{|\cdot|}(t, \Delta)$ , that  $K_1 \varepsilon^{2n/\alpha} \leq N_{\mathcal{O}_1}(\varepsilon) \leq K_2 \varepsilon^{2n/\alpha}$  for  $\varepsilon \in (0, 1]$  and  $K_1 x^n \leq N_{\mathcal{B}_x}(1) \leq K_2 x^n$  for  $x \geq x_0$ , for some  $K_1, K_2, x_0 \in (0, \infty)$ . This proves (2.1), (2.26) and (using (3.1)) (2.27).

## REFERENCES

- [1] Adler, R.J. (1990). *An Introduction to Continuity, Extrema, and Related Topics for General Gaussian Processes*. To appear in *IMS Lecture Notes*.
- [2] Adler, R.J. and Brown, L.D. (1986). Tail behaviour for suprema of empirical processes. *Ann. Probab.* **14** 1-30.
- [3] Adler, R.J. and Samorodnitsky, G. (1987). Tail behaviour for the suprema of Gaussian processes with applications to empirical processes. *Ann. Probab.* **15** 1339-1351.
- [4] Albin, J.M.P. (1990). On extremal theory for stationary processes. *Ann. Probab.* **18** 92-128.
- [5] Albin, J.M.P. (1990). On the upper and lower classes for a stationary Gaussian stochastic process. To appear in *Ann. Probab.*
- [6] Albin, J.M.P. (1990). On the upper and lower classes for a stationary stochastic process, with application to Rayleigh processes. Preprint.
- [7] Chevet, S. (1969).  $p$ -ellipsoides de  $l^q$ , exposant d'entropie, mesures cylindriques gaussiennes. *C. R. Acad. Sci. Paris Sect. A* **269** 658-660.
- [8] Dudley, R.M. (1973). Sample functions of the Gaussian process. *Ann. Probab.* **1** 66-103.
- [9] Fernique, X. (1970). Intégrabilité des vecteurs gaussiens. *C. R. Acad. Sci. Paris Sect. A* **270** 1698-1699.
- [10] Kôno, N. (1975). Asymptotic behavior of sample functions of Gaussian random fields. *J. Math. Kyoto Univ.* **15** 671-707.

- [11] Landau, H.J. and Shepp, L.A. (1971). On the supremum of a Gaussian process. *Sankhyā Ser. A* **32** 369-378.
- [12] Leadbetter, M.R., Lindgren, G. and Rootzén, H. (1983). *Extremes and Related Properties of Random Sequences and Processes*. Springer, New York.
- [13] Pathak, P.K. and Qualls, C. (1973). A law of iterated logarithm for stationary Gaussian processes. *Trans. Amer. Math. Soc.* **181** 185-193.
- [14] Qualls, C. and Watanabe, H. (1971). An asymptotic 0-1 behaviour of Gaussian processes. *Ann. Math. Statist.* **42** 2029-2035.
- [15] Qualls, C. and Watanabe, H. (1973). Asymptotic properties of Gaussian random fields. *Trans. Amer. Math. Soc.* **177** 155-171.
- [16] Samorodnitsky, G. (1990). Probability tails of Gaussian extrema. Preprint.
- [17] Sirao, T. and Nisida, T. (1952). On some asymptotic properties concerning Brownian motion. *Nagoya Math. J.* **4** 97-101.
- [18] Sudakov, N.V. (1969). Gaussian and Cauchy measures and  $\epsilon$ -entropy. *Soviet Math. Dokl.* **10** 310-313.
- [19] Watanabe, H. (1970). An asymptotic property of Gaussian processes. *Trans. Amer. Math. Soc.* **148** 233-248.
- [20] Weber, M. (1980). Analyse asymptotique des processus Gaussiens stationnaires. *Ann. Inst. H. Poincaré Sect. B* **16** 117-176.
- [21] Weber, M. (1990). The supremum of Gaussian processes with a constant variance. Preprint.

Institutionen för Matematisk Statistik  
 Lunds Universitet  
 Box 118, S-221 00 Lund, Sweden

Technical Reports  
Center for Stochastic Processes  
Department of Statistics  
University of North Carolina  
Chapel Hill, NC 27599-3260

- 258. C. Houdré, Linear Fourier and stochastic analysis, Apr. 89.
- 259. G. Kallianpur, A line grid method in areal sampling and its connection with some early work of H. Robbins, Apr. 89. *Amer. J. Math. Manag. Sci.*, 1989, to appear.
- 260. G. Kallianpur, A.G. Miamee and H. Niemi, On the prediction theory of two-parameter stationary random fields, Apr. 89. *J. Multivariate Anal.*, 32, 1990, 120-149.
- 261. I. Herbst and L. Pitt, Diffusion equation techniques in stochastic monotonicity and positive correlations, Apr. 89.
- 262. R. Selukar, On estimation of Hilbert space valued parameters, Apr. 89. (*Dissertation*)
- 263. E. Mayer-Wolf, The noncontinuity of the inverse Radon transform with an application to probability laws, Apr. 89.
- 264. D. Monrad and W. Philipp, Approximation theorems for weakly dependent random vectors and Hilbert space valued martingales, Apr. 89.
- 265. K. Benhenni and S. Cambanis, Sampling designs for estimating integrals of stochastic processes, Apr. 89.
- 266. S. Evans, Association and random measures, May 89.
- 267. H.L. Hurd, Correlation theory of almost periodically correlated processes, June 89.
- 268. O. Kallenberg, Random time change and an integral representation for marked stopping times, June 89. *Probab. Th. Rel. Fields*, accepted.
- 269. O. Kallenberg, Some uses of point processes in multiple stochastic integration, Aug. 89.
- 270. W. Wu and S. Cambanis, Conditional variance of symmetric stable variables, Sept. 89.
- 271. J. Mijneer, U-statistics and double stable integrals, Sept. 89.
- 272. O. Kallenberg, On an independence criterion for multiple Wiener integrals, Sept. 89.
- 273. G. Kallianpur, Infinite dimensional stochastic differential equations with applications, Sept. 89.
- 274. G.W. Johnson and G. Kallianpur, Homogeneous chaos, p-forms, scaling and the Feynman integral, Sept. 89.
- 275. T. Hida, A white noise theory of infinite dimensional calculus, Oct. 89.
- 276. K. Benhenni, Sample designs for estimating integrals of stochastic processes, Oct. 89. (*Dissertation*)
- 277. I. Rychlik, The two-barrier problem for continuously differentiable processes, Oct. 89.
- 278. G. Kallianpur and R. Selukar, Estimation of Hilbert space valued parameters by the method of sieves, Oct. 89.

279. G. Kallianpur and R. Selukar, Parameter estimation in linear filtering, Oct. 89.
280. P. Bloomfield and H.L. Hurd, Periodic correlation in stratospheric ozone time series, Oct. 89.
281. J.M. Anderson, J. Horowitz and L.D. Pitt, On the existence of local times: a geometric study, Jan. 90.
282. G. Lindgren and I. Rychlik, Slepian models and regression approximations in crossing and extreme value theory, Jan. 90.
283. H.L. Koul, M-estimators in linear models with long range dependent errors, Feb. 90.
284. H.L. Hurd, Almost periodically unitary stochastic processes, Feb. 90.
285. M.R. Leadbetter, On a basis for 'Peaks over Threshold' modeling, Mar. 90.
286. S. Cambanis and E. Masry, Trapezoidal stratified Monte Carlo integration, Mar. 90.
287. M. Marques and S. Cambanis, Dichotomies for certain product measures and stable processes, Mar. 90.
288. M. Maejima and Y. Morita, Trimmed sums of mixing triangular arrays with stationary rows, Mar. 90.
289. S. Cambanis and M. Maejima, Characterizations of one-sided linear fractional Lévy motions, Mar. 90.
290. N. Kono and M. Maejima, Hölder continuity of sample paths of some self-similar stable processes, Mar. 90.
291. M. Merkle, Multi-Hilbertian spaces and their duals, Mar. 90.
292. H. Rootzén, M.R. Leadbetter and L. de Haan, Tail and quantile estimation for strongly mixing stationary sequences, Apr. 90.
293. K. Benhenni and S. Cambanis, Sampling designs for estimating integrals of stochastic processes using quadratic mean derivatives, Apr. 90.
294. S. Nandagopalan, On estimating the extremal index for a class of stationary sequences, Apr. 90.
295. M.R. Leadbetter and H. Rootzén, On central limit theory for families of strongly mixing additive set functions, May 90.
296. W. Wu, E. Carlstein and S. Cambanis, Bootstrapping the sample mean for data from general distribution, May 90.
297. S. Cambanis and C. Houdré, Stable noise: moving averages vs Fourier transforms, May 90.
298. T.S. Chiang, G. Kallianpur and P. Sundar, Propagation of chaos and the McKean-Vlasov equation in duals of nuclear spaces, May 90.
299. J.M.P. Albin, On the upper and lower classes for stationary Gaussian fields on Abelian groups with a regularly varying entropy, June 90.